

On the value of coupling constant.

P.I.Pronin ^{*}and K.V.Stepanyantz [†]

February 1, 2008

*Moscow State University, physical faculty,
department of theoretical physics.
117234, Moscow, Russia*

Abstract

Using an analogy between Bogomol'nyi bound and the harmonic oscillator in quantum mechanics we propose a possible explanation of the coupling constant numerical value at the grand unification scale. It is found to be $1/8\pi$.

According to the recent precise experiments [1], the running coupling constants do not meet in a single point in the standard model. Nevertheless, in its supersymmetric extension (MSSM) [2] it really takes place. (see for example [3]). Usually, it was used as an indirect proof of supersymmetry existence. Nevertheless, this fact can be also considered from a different point, namely as an indirect measurement of α_{GUT} (the coupling constant value at the scale of Grand Unification). Numerically it appeared to be approximately $1/25$.

At the present time there is no theoretical explanation why the coupling constant has a definite value. In our opinion the Theory Of Everything should predict this value. Possibly, it is a simple combination of mathematical constants. Let us note, that numerical value of α_{GUT}^{-1} , produced by the indirect experimental data in MSSM, is very close to $8\pi \approx 25.13$. Of course, this

^{*}E-mail:pronin@theor.phys.msu.su

[†]E-mail:stepan@theor.phys.msu.su

value depends on the renormalization scheme beyond the one-loop approximation. Nevertheless, say, in the Quantum Electrodynamics the measurable values can be obtained by calculations in the definite scheme. That is why below we will suggest the possibility of α_{GUT} measurement and the existence of some preferable (physical) scheme.

In this paper we will also try to seek for a possible explanation why the coupling constant has a definite value. For this purpose we will use the analogy between the harmonic oscillator in Quantum Mechanics and the Bogomol'nyi bound.

Really, the harmonic oscillator is the simplest example of the self-dual model [4], because its Hamiltonian

$$H = \frac{p^2}{2m} + \frac{m\omega^2 x^2}{2} \quad (1)$$

is invariant under the transformations

$$p \rightarrow -m\omega x \quad x \rightarrow p/m\omega \quad (2)$$

From the other hand, first duality conjectures [5] were based on the symmetry of Dirac quantization condition and Bogomol'nyi bound [6, 4]

$$M \geq v(Q_e^2 + Q_m^2)^{1/2} \quad (3)$$

where Q_e and Q_m are electric and magnetic charges and M is a mass of state.

Let us note, that the right hand side of (3) looks very like (1).

The commutator of electric and magnetic fluxes through the closed loops α and β respectively is [7]

$$[\hat{B}(\alpha), \hat{E}(\beta)] = iGL(\alpha, \beta) \quad (4)$$

where $GL(\alpha, \beta)$ is a Gauss linking number between the loops α and β .

Assuming the commutator of electric and magnetic charges to be a limit of the commutator through the finite contours, we found, that it is not well defined. In this case we will substitute GL in (4) by an arbitrary (undefined) integer m , so that

$$[\hat{Q}_e, \hat{Q}_m] = im \quad (5)$$

The simplest way to satisfy this is to identify

$$\begin{aligned}\hat{Q}_e &= e; \\ \hat{Q}_m &= \frac{m}{i} \frac{d}{de}\end{aligned}\tag{6}$$

Then, using the analogy with the harmonic oscillator, we write down a Schrodinger type equation

$$\left(-m^2 \frac{d^2}{de^2} + e^2\right) \psi(e) = 4\pi\alpha\psi(e),\tag{7}$$

(4π comes from $\alpha_{class} = e^2/4\pi$ in the chosen unit system)

It is natural to suggest, that $\psi(e)$ is a wave function for the coupling constant distribution and eigenvalues of the operator in the left hand are observables (instead of Q_e^2 and Q_m^2). They are discrete and equal to $m(1+2n)$. The lowest nontrivial eigenvalue is 1 with the corresponding eigenfunction

$$\psi_0 = \frac{1}{\pi^{1/4}} \exp(-e^2/2).\tag{8}$$

Therefore, at the first sight $\alpha = 1/4\pi$. Nevertheless, it is valid only above the Grand Unification Scale. Really, the derivative over the coupling constant originates from magnetic charge. In Grand Unification models its mass is about M_{GUT} [8] and, therefore, the corresponding term is absent in the low energy physics. So, below the threshold we should omit the operator of squared magnetic charge and calculate the middle value of

$$\alpha_{GUT} = \frac{\langle e^2 \rangle}{4\pi} = \frac{1}{4\pi} \int de \, e^2 \frac{1}{\sqrt{\pi}} \exp(-e^2) = \frac{1}{8\pi}.\tag{9}$$

Then the numerical value of α_{GUT}^{-1} is approximately 25.13.¹, that agree well with the indirect predictions of MSSM [9], if we take into account recent experimental data [1].

As we already mentioned, above the Grand Unification scale

$$\alpha_{GUT} = \frac{1}{4\pi}.\tag{10}$$

¹It is easy to see, that the result does not depend on the particular choice of unit system.

The behavior of running coupling constants is sketched it at the Fig.1.




Figure 1: Running of the coupling constants.

To conclude we want to mention, that the main ideas of this paper are rather guesses and analogies, than results. But the hope, that something may be accidentally true, gave life to the present paper. At least the equation $\alpha_{GUT} = 1/8\pi$ seems rather curious.

References

- [1] Review of Particle Properties, *Phys.Rev.* **D50**, 1173, (1994).
- [2] For review see H.Nilles, *Phys.Rep.* **110**, 1, (1984).
- [3] K.Dienes, *Phys.Rept.*, **287**, 447, (1997).
- [4] J.Harvey, *Magnetic monopoles, duality and supersymmetry*, hep-th/9603086.

- [5] C.Montonen and D.Olive, *Phys.Lett.* **72 B**, 177, (1977).
- [6] E.B.Bogomol'nyi, *Sov.J.Nucl.Phys.*, **24**, (1976), 449.
- [7] A.Ashtekar, A.Corichi, *Phys.Rev.*, **D 56**, (1997), 2073.
- [8] C.Dokos, T.Tomaras, *Phys.Rev.* **D 21**, 2940, (1980).
- [9] S.Dimopoulos, S.Raby and F.Wilczek, *Phys.Rev.* **D24**, 1681, (1979);
S.Dimopoulos and H.Georgy, *Nucl.Phys.* **B193**, 150, (1981).